arXiv:gr-qc/0001041v3 23 Aug 2000

gr-qc/0001041
OU-TAP 117
YITP-00-16

# On cosmological perturbations in FRW model with scalar field and false vacuum decay

Arsen Khvedelidze [a,b], George Lavrelashvili [a] and Takahiro Tanaka [c,d]

[a] *Department of Theoretical Physics, A.Razmadze Mathematical Institute, GE-380093 Tbilisi, Georgia*
[b] *Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia*
[c] *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*
[d] *Department of Earth and Space Science, Graduate School of Science Osaka University, Toyonaka 560-0043, Japan*

The unconstrained reduced action corresponding to the dynamics of scalar fluctuations about the Friedmann-Robertson-Walker (FRW) background is derived using Dirac's method of description of singular Lagrangian systems. The results are applied to so-called negative mode problem in description of tunneling transitions with gravity. With our special choice of physical variable, the kinetic term of the reduced action has a conventional signature for a wide class of models. In this representation, the existence of a negative mode justifying the false vacuum decay picture turns out to be manifest. We also explain how the present result becomes consistent with the previously proved "no negative mode (supercritical supercurvature mode) theorem".

PACS number(s): 98.80.Hw, 98.80.Cq

One of the interesting and widely exploiting cosmological models is based on the theory of a scalar field coupled to gravity. Such a remarkable phenomena as inflation and metastable (false) vacuum decay are usually discussed in frame of this model. For complete description of this processes, it is very important to know the properties of perturbations to the background configurations describing an inflation or vacuum decay. The cosmological perturbations in Lorentzian regime are related to the cosmic microwave background radiation and large scale structure formation [1–3]. Furthermore the perturbations in the Euclidean version of the theory define one loop corrections to the bubble nucleation rate and determine quantum state of materialized bubble [4]. Since the model contains gauge degrees of freedom, the choice of physical variables to analyze the system is not unique.

There are several known ways in deriving and expressing unconstrained quadratic action solely written in terms of physical variables in the theory of scalar field coupled to gravity in non-spatially flat FRW Universe [2,3,5–10]. However, none of them is completely satisfactory for the purpose to apply it to the issue of quntum tunneling, which we discuss mainly in this short letter. An extension of the convenient variable $v$ discussed in the well-known review [1] to the non-spatially flat case is found in Ref. [10]. However, the canonical transformation to arrive at this variable turns out to be singular in the case of quantum tunneling. In the reduction discussed in [5], the kinetic term does not have definite signature. In the case of the reduction given in [6,2], the kinetic term has a definite signature (as long as the background scalar field is monotonic). However, the overall signature is unconventional and hence some analytic continuation similar to the conformal rotation becomes necessary. The conformal rotation was introduced in the case of pure gravity to cure the conformal factor problem [11]. It was also found to be possible to treat this problem (at least in perturbative gravity) via the careful gauge fixing [12].

In this short letter, we revisit the dynamics of small scalar-type perturbations of a scalar field coupled to FRW type background in the framework of conventional theory of degenerate Lagrangian systems [13]. We obtain an expression for the reduced action which is more appropriate for discussing the quantum tunneling with gravity. For a wide class of models, the signature of the obtained reduced action becomes the conventional one. The potential for the eigen value problem is also regular for the same class of models, and it is shown that there is a negative mode in the spectrum of small fluctuations about the Coleman-De Luccia bounce solution [15] in accordance with the consistent interpretation of metastable vacuum decay [4]. As it will become clear later, this result does not conflict with the conclusions about the absence of negative mode arrived in the other reduction scheme [2,6–9].

The evolution of system composed of scalar matter field minimally coupled to gravity is determined by the conventional action

$$S = \int d^4x\sqrt{-g}\left[\frac{1}{2\kappa}R - \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi - V(\phi)\right] , \qquad (1)$$

where $\kappa = 8\pi G$ is the reduced Newton's constant and $V(\phi)$ is the scalar field potential.

We expand the metric and the scalar field over an FRW type background

$$\begin{aligned}ds^2 = a(\eta)^2 \Bigg[ &-(1+2A(\eta)Y)d\eta^2 + 2\mathcal{B}(\eta)Y_{|i}d\eta dx^i \\ &+ \left\{\gamma_{ij}(1-2\Psi(\eta)Y) + 2\mathcal{E}(\eta)Y_{|ij}\right\} dx^i dx^j \Bigg], \\ \phi = \varphi(\eta) &+ \Phi(\eta)Y, \end{aligned} \tag{2}$$

where $\gamma_{ij}$ is the three-dimensional metric on the constant curvature space sections, $a$ and $\varphi$ are the background field values and $A, \Psi$, $\Phi$, $\mathcal{B}$ and $\mathcal{E}$ are small perturbations. $Y$ is a normalized eigen function of 3-dimensional Laplacian, $\Delta Y = -k^2 Y$, and vertical line represents a covariant derivative with respect to $\gamma_{ij}$. To keep the simplicity, we set $\mathcal{B}(\eta) = \mathcal{E}(\eta) = 0$. These terms are absent for homogeneous perturbations from the beginning. Also for inhomogeneous perturbations, it is known that this choice of gauge is consistent.

The background fields $a$ and $\varphi$ satisfy the equations

$$\mathcal{H}^2 - \mathcal{H}' + \mathcal{K} = \frac{\kappa}{2}\,\varphi'^2, \tag{3}$$

$$2\,\mathcal{H}' + \mathcal{H}^2 + \mathcal{K} = \frac{\kappa}{2}\left(-\varphi'^2 + 2\,a^2 V(\varphi)\right), \tag{4}$$

$$\varphi'' + 2\,\mathcal{H}\varphi' + a^2\frac{\delta V}{\delta\varphi} = 0, \tag{5}$$

where a prime denotes a derivative with respect to conformal time $\eta$, $\mathcal{H} := a'/a$, and $\mathcal{K}$ is the curvature parameter, which has the values 1, 0, -1 for closed, flat and open universes, respectively.

Expanding the total action, keeping terms of second order in perturbations, and using the background equations, we find

$$S = S^{(0)} + S^{(2)}, \tag{6}$$

where $S^{(0)}$ is the action for the background solution and $S^{(2)}$ is quadratic in perturbations with the Lagrangian for scalar perturbations

$$\begin{aligned}{}^{(s)}\mathcal{L} = \frac{a^2\sqrt{\gamma}}{2\kappa}\Bigg[ &-6\Psi'^2 + 2(k^2 - 3\mathcal{K})\Psi^2 \\ &+\kappa\left\{\Phi'^2 - \left(a^2\frac{\delta^2 V}{\delta\varphi\delta\varphi} + k^2\right)\Phi^2 + 6\varphi'\Psi'\Phi\right\} \\ &-\Big\{2\kappa\varphi'\Phi' + 2\kappa a^2\frac{\delta V}{\delta\varphi}\Phi \\ &\qquad +12\mathcal{H}\Psi' + 4(k^2 - 3\mathcal{K})\Psi\Big\}A \\ &-2(\mathcal{H}' + 2\mathcal{H}^2 - \mathcal{K})A^2\Bigg]. \end{aligned} \tag{7}$$

To obtain the unconstrained system corresponding to the degenerate Lagrangian (7) we will follow the Dirac's description of singular Lagrangian systems [13]. Performing the Legendre transformation with canonical momenta

$$\Pi_\Psi := \frac{\delta\,{}^{(s)}\mathcal{L}}{\delta\Psi'} = \frac{6\,a^2\sqrt{\gamma}}{\kappa}\left(-\Psi' + \frac{\kappa}{2}\varphi'\Phi - \mathcal{H}\,A\right), \tag{8}$$

$$\Pi_\Phi := \frac{\delta\,{}^{(s)}\mathcal{L}}{\delta\Phi'} = a^2\sqrt{\gamma}\,(\Phi' - \varphi'\,A), \tag{9}$$

$$\Pi_A := \frac{\delta\,{}^{(s)}\mathcal{L}}{\delta A'} = 0, \tag{10}$$

we find the primary constraint $C_1 := \Pi_A = 0$. Thus the total Hamiltonian $H_T$ is

$$H_T = H_C + u_1(\eta)C_1, \tag{11}$$

with arbitrary function $u_1(\eta)$ and canonical Hamiltonian

$$\begin{aligned}H_C = &-\frac{\kappa}{12\,a^2\sqrt{\gamma}}\Pi_\Psi^2 + \frac{1}{2\,a^2\sqrt{\gamma}}\Pi_\Phi^2 + \frac{\kappa}{2}\varphi'\,\Pi_\Psi\Phi \\ &+a^2\sqrt{\gamma}\Bigg[-\frac{k^2 - 3\mathcal{K}}{\kappa}\Psi^2 \\ &+\frac{1}{2}\left(a^2\frac{\delta^2 V}{\delta\varphi\delta\varphi} - \frac{3}{2}\kappa\varphi'^2 + k^2\right)\Phi^2\Bigg] + A\,C_2\,, \end{aligned} \tag{12}$$

where

$$\begin{aligned}C_2 = &\varphi'\Pi_\Phi - \mathcal{H}\Pi_\Psi \\ &+ a^2\sqrt{\gamma}\left[\left(a^2\frac{\delta V}{\delta\varphi} + 3\varphi'\mathcal{H}\right)\Phi + \frac{2(k^2 - 3\mathcal{K})}{\kappa}\Psi\right]. \end{aligned} \tag{13}$$

Conservation of primary constraint gives the secondary constraint $C_2 = 0$. The primary and the secondary constraints are first class and there are no ternary constraints.

The existence of constraints in the system as usually means the presence of gauge degrees of freedom. To identify the physical degrees of freedom, we fix the gauge and solve the constraints. There are two simple strategies: either to eliminate perturbations of scalar field ($\Pi_\Phi$ and $\Phi$) or the gravitational degrees of freedom ($\Pi_\Psi$ and $\Psi$). The approach developed in [6–9,2] is based on the first possibility.[*] Here we use the second possibility.

Thus we choose the following gauge fixing condition

$$\chi_1 := \Pi_\Psi = 0\,. \tag{14}$$

From the consistency condition $\chi_1' = 0$, we obtain

$$\chi_2 = A - \Psi = 0, \tag{15}$$

which is the condition known as the Newton gauge. Then, the consistency condition $\Psi' - A' = 0$ will determine $u_1$, and the set of constraints closes.

As a next step one can introduce the Dirac brackets, or equivalently we can identify the Hamiltonian for the physical degrees of freedom $H^*$ by the relation

---

[*] Note that the gauge invariant formulation in [2] corresponds to the gauge choice $\chi_{GMST} := \Phi = 0$.

$$\Pi_\Psi \Psi' + \Pi_\Phi \Phi' + \Pi_A A' - H_C\big|_{\chi_i=0, C_i=0} = \Pi_\Phi \Phi' - H^* . \tag{16}$$

After some algebra, assuming $k^2 \neq 3\mathcal{K}$, we obtain

$$\begin{aligned} H^* = {} & \frac{Q}{2a^2\sqrt{\gamma}}\Pi_\Phi^2 - \frac{\kappa\varphi'}{2(k^2-3\mathcal{K})}\left(3\varphi'\mathcal{H} + a^2\frac{\delta V}{\delta\varphi}\right)\Pi_\Phi\Phi \\ & + \frac{1}{2}a^2\sqrt{\gamma}\Bigg[-\frac{\kappa}{2(k^2-3\mathcal{K})}\left(3\varphi'\mathcal{H} + a^2\frac{\delta V}{\delta\varphi}\right)^2 \\ & \qquad + \left(a^2\frac{\delta^2 V}{\delta\varphi\delta\varphi} - \frac{3}{2}\kappa\varphi'^2 + k^2\right)\Bigg]\Phi^2, \end{aligned} \tag{17}$$

with

$$Q := 1 - \frac{\kappa\varphi'^2}{2(k^2-3\mathcal{K})} \, . \tag{18}$$

To proceed further it is useful to introduce new canonical coordinates $f$ and $\pi_f$ defined by

$$\begin{aligned} \pi_f & - \left[Q\mathcal{H} + \frac{\kappa\varphi'}{6\mathcal{K}}\frac{1+Q}{Q}\left(2\varphi'\mathcal{H} + a^2\frac{\delta V}{\delta\varphi}\right)\right] f \\ & = \sqrt{\frac{Q}{a^2\sqrt{\gamma}}}\Pi_\Phi, \end{aligned} \tag{19}$$

$$f = \sqrt{\frac{a^2\sqrt{\gamma}}{Q}}\Phi. \tag{20}$$

One finds that the dynamics of physical variable $f$ is governed by the simple harmonic oscillator Hamiltonian

$$H^* = \frac{1}{2}\pi_f^2 + \frac{1}{2}w^2[a(\eta), \varphi(\eta)] f^2, \tag{21}$$

with frequency whose time dependence is determined by the background solutions

$$w^2[a(\eta),\varphi(\eta)] = \frac{\sqrt{Q}}{\varphi'}\left(\frac{\varphi'}{\sqrt{Q}}\right)'' - (k^2-3\mathcal{K})Q. \tag{22}$$

Thus, the unconstrained quadratic action becomes

$$S^{(2)} = \int\left(\frac{1}{2}f'^2 - \frac{1}{2}w^2[a(\eta),\varphi(\eta)]f^2\right)d\eta \, . \tag{23}$$

Now we apply the derived reduced action to investigation of the negative mode problem in quantum tunneling with gravity. The false vacuum decay is described by the bounce solution of Euclidean equations [14,15]. Value of the Euclidean action at the bounce gives leading exponential factor in decay rate. Quadratic action defines one loop corrections. It is remarkable that in the spectrum of small perturbations about a bounce in absence of gravity there is exactly one negative mode [4]. This mode is responsible for making correction to the ground-state energy imaginary, i.e., justifying decay interpretation.

Relevant object for tunneling transitions is the Euclidean action which can be obtained from the action Eq. (23) by the analytic continuation $\eta = -i\tau$. Defining the Euclidean action as usual by $S^{(2)} = iS_E^{(2)}$ and specifying $\mathcal{K} = +1$, we obtain

$$S_E^{(2)} = 2\pi^2\int\left(\frac{1}{2}\dot{f}^2 + \frac{1}{2}U[a(\tau),\varphi(\tau)]f^2\right)d\tau \, , \tag{24}$$

where $\dot{} = d/d\tau$ and

$$U = \frac{\sqrt{Q_E}}{\dot\varphi}\left(\frac{d^2}{d\tau^2}\frac{\dot\varphi}{\sqrt{Q_E}}\right) + (\ell(\ell+2)-3)Q_E, \tag{25}$$

with

$$Q_E = 1 + \frac{\kappa\dot\varphi^2}{2(\ell(\ell+2)-3)}. \tag{26}$$

Here we used the fact that the eigen values of the Laplacian $\Delta$ on a unit sphere take discrete values, i.e., $k^2 = \ell(\ell+2)$ with $\ell = 0, 1, 2, \ldots$.

The equation for the mode functions, which diagonalize the action Eq.(24), has form of the Schrödinger equation

$$-\frac{d^2}{d\tau^2}f + U[a(\tau),\varphi(\tau)]\, f = E\, f \, . \tag{27}$$

Let's us first show that for $\ell = 0$ case the Eq.(27) has at least one negative mode for the Coleman-De Luccia background bounce solution. We define a new potential

$$\tilde{U} := \frac{\sqrt{Q_E}}{\dot\varphi}\left(\frac{d^2}{d\tau^2}\frac{\dot\varphi}{\sqrt{Q_E}}\right) > U. \tag{28}$$

The eigen value problem with this potential manifestly has a zero eigen value state with $f \propto \dot\varphi/\sqrt{Q_E}$. Since $\tilde{U} > U$, the eigen value problem with the potential $U$ must have at least one negative eigen mode. □

A similar discussion leads to the conclusion that there are no negative modes for $\ell > 1$ (compare [7]). The $\ell = 1$ case needs separate consideration. As it was explained in Ref. [6], there are no physical degrees of freedom in this sector.

Note that the present result does not contradict with the no negative mode theorem [9]. The argument of the no negative mode theorem is that, when we consider the bounce solution such that realizes the minimum value of action among all the non-trivial $O(4)$-symmetric configurations, there is no negative mode for the specific variable $q$ defined by

$$q := \frac{8a}{\sqrt{\kappa}\dot\varphi}\Psi^N, \tag{29}$$

where $\Psi^N$ represents $\Psi$ evaluated in the Newton gauge [2]. In terms of this variable $q$, the kinetic term stays negative for $\ell = 0$ modes. Hence, as mentioned earlier, we need to do some analytic continuation similar to the conformal rotation to perform the path integral. Although it is not fully justified, this procedure of analytic continuation is thought to produce the required imaginary factor. (See. [6,9]). Now we are working with a different variable $f$ for which the kinetic term takes the conventional signature. Therefore there should be one and only one negative mode.

Once we accept the original no negative mode theorem, we can give an indirect proof of the uniqueness of negative mode for $f$ under the same condition that the absence of negative mode for $q$ was proven. It is easy to find the relation between $f$ and $q$ because our gauge condition corresponds to the Newton gauge (15). Eliminating $\Pi_\Phi$ from the constraint equation $C_2 = 0$ and the Hamiltonian equation of motion for $\Phi$ derived from (17), we obtain

$$\frac{d}{d\tau}\left(\frac{\sqrt{Q_E}}{\sqrt[4]{\gamma}\dot{\varphi}} f\right) = \frac{\ell(\ell+2) - 3\mathcal{K}}{4\sqrt{\kappa}\dot{\varphi}} Q_E\, q, \tag{30}$$

where we used the definitions (20) and (29). Suppose that $f$ has two negative modes for $\ell = 0$, and let us try to derive a contradiction by using the fact that $q$ doesn't have any negative mode. By assumption, the zero eigen value solution ($E = 0$) of $f$ which satisfies the boundary condition on one side must have two nodes. One may think that this implies that there are at least two zeros of $z := (\sqrt{Q_E}/\sqrt[4]{\gamma}\dot{\varphi})f$. But it is not true because the regularity condition for $f$ just requires it to behave like $e^{-(\ell+1)|\tau|}$ on boundaries. Since $\dot{\varphi}$ behaves like $\sim e^{-2|\tau|}$, even the regular solution of $z$ does not go to zero on the boundary for $\ell = 0$. Hence, we arrive at the conclusion that there is at least one point where the derivative of $z$ vanishes. With the aid of Eq.(30), this implies that the zero eigen value solution of $q$ has a node, which means existence of a negative mode in $q$ and contradicts with the no-negative mode theorem. □

We also note that with the present choice of variable there is no manifest correspondence between no negative mode theorem and the no supercritical supercurvature mode theorem [8] as was in terms of variable $q$ defined in [2]. The equation which determines the perturbation spectrum in the context of one bubble open inflation in terms of the present variable $f$ is not standard Schrödinger-type equation and the existence of a mode with a negative value of $E$ does not imply the existence of a supercritical supercurvature mode.

So far we considered the background solutions with the positive definite factor $Q_E$. If this factor becomes zero or negative for some region(s) of $\tau$, the Euclidean analog of the canonical transformation Eqs.(19), (20) becomes singular. This will not immediately indicate a certain physical meaning because there happens nothing special as long as we discuss in terms of $q$ in Ref. [2]. However, if there exist some class of bounce solutions for which the signature of kinetic term cannot be set to be positive definite without passing through a singular canonical transformation, then it might suggest some physical meaning. In this case, we might have to reconsider the possibility of catastrophic particle creation discussed in Ref. [5].

To conclude we have investigated the dynamics of small perturbations in non-flat FRW model coupled to a scalar field. Using the gauge conditions Eq.(14) we reduced the system of coupled perturbations Eq.(7) to the dynamical system Eq.(23) with one physical degree of freedom. The reduced quadratic action Eq.(23) has the conventional overall signature. Investigating Euclidean quadratic action Eq.(24), we proved that there is exactly one negative eigen value mode about the Coleman-De Luccia bounce solution. The result is consistent with the so-called "no-negative mode" theorem [9] in false vacuum decay with gravity which was proven in the other reduction scheme. The treatment discussed here is restricted to the background solutions which satisfy the condition that the quantity $Q_E$ is positive. Hence, another question arises whether we can always find a variable for which the kinetic term of a perturbation in the reduced action becomes positive definite. This issue needs further investigation.

## ACKNOWLEDGMENTS

G.L. is grateful to D. Maison, V.A. Rubakov and G. Tsitsishvili for many interesting discussions and comments at different stages of this work. T.T. thanks J. Garriga and X. Montes for valuable comments. Work of G.L. was partly supported by the Georgian Academy of Sciences under Grant No 1.4 and by the Royal Society Grant. Work of T.T. was partly supported by Monbusho System to Send Japanese Researchers Overseas.

---

[1] V.F. Mukhanov, H.A. Feldmann and R.H. Brandenberger, *Theory of cosmological perturbations*, *Phys. Rep.* **215**, (1992), 203-333.

[2] J. Garriga, X. Montes, M. Sasaki and T. Tanaka, *Canonical Quantization of Cosmological Perturbations in the One Bubble Open Universe*, *Nucl. Phys.* **B513**, (1998), 343-374.

[3] S. Gratton and N. Turok, *Cosmological Perturbations from the No Boundary Euclidean Path Integral*, *Phys. Rev.* **60**, (1999), 123507.

[4] C.G. Callan and S. Coleman, *Fate of the false vacuum. II. First quantum corrections*, *Phys. Rev.* **D16**, (1977), 1762-1768.

[5] G.V. Lavrelashvili, V.A. Rubakov and P.G. Tinyakov, *Tunneling transitions with gravitation: breakdown of the quasiclassical approximation*, *Phys. Lett.* **161B**, (1985), 280-284.

[6] T. Tanaka and M. Sasaki, *False Vacuum Decay with Gravity. Negative Mode Problem*, *Prog. of Theor. Phys.* **88**, (1992), 503-528.

[7] G. Lavrelashvili, *Quadratic action of the Hawking-Turok instanton*, *Phys. Rev.* **D58**, (1998), 063505.

[8] T. Tanaka and M. Sasaki, *No supercritical supercurvature mode conjecture in one-bubble open inflation*, *Phys. Rev.* **D59**, (1999), 023506 , (1999).

[9] T. Tanaka, *The No-Negative Mode Theorem in False Vacuum Decay with Gravity*, *Nucl. Phys.* **B556**, (1999), 373-396.

[10] J. Garriga, and V.F. Mukhanov, *Perturbations in k-inflation*, *Nucl. Phys.* **B458**, (1999), 219-225.

[11] G.W. Gibbons, S.W. Hawking and M.J. Perry, *Path integrals and the indefiniteness of the gravitational action*, *Nucl. Phys.* **B138**, (1978) 141-150.

[12] J. Hartle and K. Schleich, The conformal rotation in linearized gravity, in Quantum Field Theory and Quantum Statistics, **v2**, 67-87, Eds. T.A. Batalin, C.J. Isham, and G.A. Vilkovisky, Hilger, Bristol, 1987.

[13] P.A.M. Dirac, *Lectures on Quantum Mechanics.* Belfer Graduate School of Science, (Yeshiva University, New York, 1964).

[14] S. Coleman, *Fate of the false vacuum: Semiclassical Theory*, *Phys. Rev.* **D15**, (1977), 2929-2936.

[15] S. Coleman and F. De Luccia, *Gravitational effects on and of vacuum decay*, *Phys. Rev.* **D21**, 3305-3313, (1980).